

FluSure XP/RespiSure®

Pfizer Animal Health

Swine Influenza Vaccine, H1N1, H1N2 & H3N2, Killed Virus and
Mycoplasma Hyopneumoniae Bacterin

INDICATIONS

For vaccination of healthy swine, including pregnant sows and gilts, 3 weeks of age or older as an aid in preventing respiratory disease caused by SIV subtypes H1N1, H1N2 and H3N2 and *M. hyopneumoniae*. A duration of immunity of at least 10 weeks, as demonstrated by a significant reduction in lung lesions, has been established against SIV H3N2. This product contains the following SIV isolates: A/Swine/North Carolina/031/05 (H1N1); A/Swine/Iowa/110600/00 (H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2); and A/Swine/Missouri/069/05 (H3N2).

DIRECTIONS

Shake diluent before use. Aseptically rehydrate the freeze-dried vaccine with the liquid bacterin provided, shake well, and administer 2 mL intramuscularly. Healthy swine 3 weeks of age or older should receive two 2-mL doses administered approximately 3 weeks apart. In young pigs, vaccinate after maternally derived antibodies to SIV have declined. Semiannual revaccination with a single dose is recommended.

PRECAUTIONS

Store at 2°–7°C. Do not freeze. Use entire contents when first opened or rehydrated. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Contains gentamicin as preservative. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

Technical inquiries should be directed to Pfizer Animal Health Veterinary Services, (800) 366-5288 (USA), (800) 461-0917 (Canada).

DOSES

50-dose vial of vaccine, rehydrate to 100 mL
100-mL vial of liquid bacterin for use as diluent