

# COMMITTED TO QUALITY. DEDICATED TO ANIMAL HEALTH.

## Cocci Cure® C for Calves

(amprolium)



### PRODUCT DESCRIPTION:

An oral solution used as aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. Medicated drinking water is palatable and readily accepted by calves.

- Breaks up the Coccidia life cycle
- Approved for the prevention and treatment of Coccidiosis in beef and dairy cattle
- Medicated drinking water is palatable and readily accepted by calves
- One gallon of concentrate treats 1,600 gallons of drinking water based on 21-day prevention regimen
- 24-hour slaughter withdrawal time after the last administration
- Treat the entire herd through the drinking water
- Concentrated solution disperses readily into the drinking water
- Approved by FDA

### SIZES:

1 G (128 fl oz) (3.785 L)

Store container in an upright position. Store between 5°C-25°C (41°F-77°F), with brief excursions to 40°C.

See package insert or label for complete indications, precautions, warnings, residue information, and use directions.



# Cocci Cure® C for Calves

(amprolium) 9.6% Oral Solution Coccidiostat

## PACKAGING LABEL:

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**  
**USE DIRECTIONS:** There are **96 mg** of amprolium in every **1 mL** of Cocci-Cure® C FOR CALVES.

**5-Day Treatment Protocol**  
Daily Dosage: 10 mg amprolium/kg  
(10 mg per 2.2 lb body weight)

**21-Day Prevention Protocol**  
Daily Dosage: 5 mg amprolium/kg  
(5 mg per 2.2 lb body weight)

1 fl oz = 29.57 mL  
1 fl oz = 2 measuring tablespoons  
8 fl oz = 1/2 pint (236.56 mL)  
16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER:** Mix Cocci-Cure® C FOR CALVES daily with fresh drinking water.

**5-DAY TREATMENT:** Add Cocci-Cure® C FOR CALVES to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add Cocci-Cure® C FOR CALVES to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

Lot No.:                      Exp. Date:



## Cocci-Cure® C FOR CALVES

**(amprolium)  
9.6% Oral Solution  
Coccidiostat**

Active Ingredient: amprolium.....9.6%

**FOR ORAL USE IN ANIMALS ONLY**

**Keep This and All Drugs  
Out of Reach of Children**

**Restricted Drug (California) - Use Only as Directed  
ANADA 200-482, Approved by FDA**

**NET CONTENTS:**  
**1 Gallon (128 fl oz) (3.785 L)**

**AS A DRENCH:** Cocci-Cure® C FOR CALVES drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz Cocci-Cure® C FOR CALVES to 1 pt (16 oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of Cocci-Cure® C FOR CALVES to 1 pt (16 fl oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNING:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: Not for human use. Keep this and all drugs out of reach of children.**

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS contact Aspen Veterinary Resources at (816.415.4324).

**PRECAUTIONS: MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention. **Sodium Benzoate 0.1% added as preservative.**

**STORAGE:** Store container in an upright position. Store between 5°C-25°C (41°F-77°F), with brief excursions to 40°C.

Cocci-Cure® is a registered trademark of Aspen Veterinary Resources, Ltd.

**TAKE TIME  OBSERVE LABEL DIRECTIONS**

Manufactured for:  
Aspen Veterinary Resources, Ltd.  
Liberty, MO 64068, USA  
www.aspenveterinaryresources.com  
A147B  
Product No.: 1AMP026  
8AMP025 Rev: 03/15



0 99355 00434 4

| Product Code | Product Size | Case   |
|--------------|--------------|--------|
| 16144354     | 1 G          | 4/Case |